

hep-ph/9807566
IEKP-KA/98-15
July 1998

Aspects of Higgs Boson Searches at LEP2

André Sopczak

University of Karlsruhe

Abstract

A preliminary combined mass limit of about 90 GeV has been reported for the Minimal Standard Model Higgs boson from the 183 GeV data taken in 1997, with a total luminosity of about 200 pb^{-1} . Similar mass limits have been reported from the individual LEP experiments based on the first 1998 data at 189 GeV with up to 40 pb^{-1} each. For the first time the LEP experiments have made a common effort to derive preliminary combined limits in the Minimal Supersymmetric extension of the Standard Model. For three MSSM benchmark parameter sets, mass limits for the lighter scalar of 77 GeV and for the pseudoscalar of 78 GeV have been derived. In a more general parameter scan unexcluded mass regions have been revealed. An independent variation of the SUSY parameters results in cancellation effects of production cross sections and thus some parameter regions are not excluded in this more general framework. Direct and indirect limits from other SUSY particle searches have been taken into account.

Presented at the Sixth International Symposium on Particles, Strings, and Cosmology,
PASCOS-98, Boston, March 22–29, 1998

ASPECTS OF HIGGS BOSON SEARCHES AT LEP2

ANDRE SOPCZAK

Karlsruhe University

E-mail: andre.sopczak@cern.ch

A preliminary combined mass limit of about 90 GeV has been reported for the Minimal Standard Model Higgs boson from the 183 GeV data taken in 1997, with a total luminosity of about 200 pb^{-1} . Similar mass limits have been reported from the individual LEP experiments based on the first 1998 data at 189 GeV with up to 40 pb^{-1} each. For the first time the LEP experiments have made a common effort to derive preliminary combined limits in the Minimal Supersymmetric extension of the Standard Model. For three MSSM benchmark parameter sets, mass limits for the lighter scalar of 77 GeV and for the pseudoscalar of 78 GeV have been derived. In a more general parameter scan unexcluded mass regions have been revealed. An independent variation of the SUSY parameters results in cancellation effects of production cross sections and thus some parameter regions are not excluded in this more general framework. Direct and indirect limits from other SUSY particle searches have been taken into account.

1 Introduction

Over the last eight years the LEP experiments have made great progress in their search for Higgs bosons. So far, despite immense efforts, no Higgs bosons have been found, and the quest to understand mass generation continues. The very successful operation of the LEP accelerator at the Z resonance until 1995 (LEP1) has continued with the successive energy increase (LEP2) to 189 GeV in 1998, and to about 200 GeV over the next two years. LEP2 operation has permitted a large sensitivity increase which has so far manifested itself in larger mass limits. In the Minimal Standard Model (MSM), mass generation results from one complex Higgs doublet. The combined LEP1 mass limit of about 65 GeV¹ has been increased to the current limit of about 90 GeV².

This article is based on the results from the four LEP experiments ALEPH, DELPHI, L3 and OPAL, and focuses on the combined limits and the interpretation in the Minimal Supersymmetric extension of the Standard Model (MSSM). The large interest in the search for Higgs bosons in the framework of the MSSM results from the fact that the MSSM predicts a light Higgs boson which could be discovered at LEP as a first sign of Supersymmetry.

2 Status of the Higgs Boson Searches

The search for the Higgs bosons has been performed for many different signatures. Each signature results from a possible Higgs boson decay mode. For

Higgs boson production, the MSM process $e^+e^- \rightarrow hZ$ gives a large cross section if the Z is produced on-shell for sufficiently large center-of-mass energy. In the MSSM, which predicts five physical Higgs bosons, the production process $e^+e^- \rightarrow hA$ can also occur. The subsequent decay of the Z and Higgs bosons determines the search channel. Most important are the Higgs boson decays into $\tau^+\tau^-$ and $b\bar{b}$. Four conceptual different statistical methods give consistent limits when the results of the four LEP experiments are combined^{3,2}. The individual limits for each experiment are given in Table 1 from ref.² The lowest combined mass limit of the four methods is 89.8 GeV, whilst the corresponding expected limit is 90.4 GeV. The spread of the observed limits from the four methods is only 0.15 GeV. The confidence level (CL) curve using the DELPHI method is given in Fig. 1. Very preliminary results from the first 1998 data-taking with up to 40 pb⁻¹ per experiment already give similar mass limits⁴ as for the total 1997 luminosity of about 200 pb⁻¹. With the combined 1998 data, a Higgs boson of about 95 GeV could be discovered, and by the end of the LEP2 programme in 2000, a Higgs boson with a mass of up to 105 GeV could be found.

Figure 1 also gives an interpretation in the MSSM of the combined 1997 data for a fixed set of SUSY parameters characterized by a large Supersymmetry scale of 1 TeV, and maximal mixing in the scalar top sector with $A = \sqrt{6}$. The exclusion confidence level has been calculated with the OPAL statistical method and mass limits of about 77 and 78 GeV are derived for scalar and pseudoscalar Higgs bosons, respectively.


Table 1: Preliminary number of data and background events in the 183 GeV data, as well as the expected and observed mass limits for the MSM Higgs boson at 95% CL. Note that when deriving the ALEPH limit, no background subtraction was applied. In the L3 analysis, the expected background and observed number of events depend on the Higgs mass hypothesis; they are given here for $m_H=85$ GeV.

Experiment	ALEPH	DELPHI	L3	OPAL
Expected background	7.2	6.5	24.2	8.9
Observed events	7	6	18	9
Expected limit (GeV)	85.5	86.5	85.0	86.2
Observed Limit (GeV)	87.9	85.7	87.6	88.3

3 MSSM Parameter Scan

The importance of a parameter scan was pointed out for LEP1⁵ and more detailed studies were performed when the LEP2 data became available⁶. New scan results from ALEPH⁷, DELPHI⁸ and OPAL⁹ have been released, and all parameter scans are based on a similar variation of the SUSY parameters.

Figure 1: Left: Preliminary MSM expected (dashed lines) and observed (solid lines) confidence levels, $CL_s(m_H)$, obtained from combining the results of the four LEP collaborations using the DELPHI statistical methods. The intersections of the curves with the 5% horizontal lines define the 95% CL lower bounds, expected and observed, for the mass of the SM Higgs boson. Right: Preliminary MSSM benchmark exclusion plot for (m_h, m_A) at 95% CL using the OPAL method to combine the different LEP experiments for $\tan\beta > 0.7$.




Details of the DELPHI scan are given as an example. The DELPHI scan includes the Higgs and SUSY particle searches of the 183 GeV data. A given (m_h, m_A) combination is excluded if, for all SUSY parameter sets from the ranges defined in Table 2, the exclusion confidence level is larger than 95% in at least one of the production reactions. The parameters shown in Table 2 are the input parameters for the calculation of the physical sfermion, chargino, and neutralino masses. Some parameter combinations can be unphysical (e.g. negative squark masses) or experimentally excluded. Such cases are removed by imposing constraints on neutralino and chargino particles¹⁰. For each parameter combination of the scan, the exclusion confidence level is calculated. In addition, the limits from the reaction $b \rightarrow s\gamma$ and the constraints on $\Delta\rho$ from light scalar top quarks¹¹ are taken into account. Figure 2 shows the excluded regions in the (m_h, m_A) plane for the preliminary DELPHI results⁸ and for an approximate combination of all four experiments based on ref⁶ OPAL obtains similar results with their parameter scan⁹. The ALEPH parameter scan⁷ points out that only a small fraction of the parameter combinations in the MSSM is not excluded compared to the benchmark results.

Table 2: Ranges of SUSY parameters used for independent variation in the study of the MSSM neutral Higgs boson searches.

Parameter	m_{sq} (GeV)	m_g (GeV)	μ (GeV)	A	$\tan\beta$
Range	200—1000	200—1000	−500—500	−1—+1	0.5—50

Figure 2: Left: Preliminary DELPHI MSSM exclusion for $\sqrt{s} = 183$ GeV and $\mathcal{L} = 54 \text{ pb}^{-1}$. The region excluded by LEP1 (black), the newly 95% CL excluded region at LEP2 (grey), the region where the exclusion depends on the SUSY parameter set (white with dots), the region with no sensitivity (white), and the theoretically not allowed region (black) are shown. Right: Approximate combined LEP MSSM exclusion for $\mathcal{L} = 200 \text{ pb}^{-1}$.



The DELPHI scan over the SUSY parameter space gives a limit of about 67 GeV on the scalar Higgs boson mass⁸ compared with a limit of 74.4 GeV for the three sets of SUSY benchmark parameter sets¹². The existence of newly unexcluded regions can be understood in the following way: The unexcluded region begins near the $m_h + m_A = \sqrt{s}$ limit for $m_h > 67$ GeV. In this range, the bremsstrahlung cross section $e^+e^- \rightarrow hZ$ can be small for some SUSY parameters. The complementary process $e^+e^- \rightarrow hA$ is reduced kinematically, and thus no signal can be observed.

The 1997 combined LEP data of about 200 pb^{-1} leads to mass limits of about 76 and 83 GeV for scalar and pseudoscalar Higgs bosons, respectively. Despite the different scan method and the additional direct and indirect limits from other SUSY particle searches applied in Fig. 2, very similar mass limits are obtained compared to the benchmark limits in Fig. 1. As pointed out in ref. ⁶, the combined luminosity of the four LEP experiments is important for a larger coverage of the MSSM parameter space. In the example of Fig. 2, the limit on the scalar Higgs boson mass increased by 9 GeV.

Acknowledgments

I would like to thank the organizers of the conference for their kind hospitality, and Wim de Boer for his advice about the manuscript.

References

1. For example: A. Sopczak, *Higgs Bosons Searches at LEP*, DESY97-129, ISSN 0418-9833, Proc. XII Int. Symposium on High Energy Physics, Gauhati, India, Dec. 1996 – Jan. 1997.
2. The LEP working group for Higgs boson searches, P. Bock, et. al., *Bounds on SM and neutral MSSM Higgs boson masses using combined data of the four LEP experiments collected at $\sqrt{s}$ up to 183 GeV*, ALEPH 98-069, PHYSIC 98-028, DELPHI 98-144, PHYS 790, L3 note 2310, OPAL Technical Note TN-558, July 17, 1998; contributed paper 582, ICHEP-98, 23–28 July, 1998, Vancouver, Canada.
3. ALEPH, DELPHI, L3 and OPAL Collaborations, The LEP working group for Higgs boson searches, *Lower bound for the Standard Model Higgs boson mass from combining the results of the four LEP experiments*, CERN-EP/98-046, April 1, 1998.
4. D. Treille, *Searching for New Particles at Existing Accelerators*, Talk ICHEP-98, 23–28 July, 1998, Vancouver, Canada.
5. J. Rosiek and A. Sopczak, *Present and future searches with e^+e^- colliders for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model – the complete 1-loop analysis*, Phys. Lett. **B341** (1995) 419.
6. A. Sopczak, *Searches for Higgs Bosons at LEP2*, hep-ph/9712283, Proc. First Int. Workshop on Non-Accelerator Physics, Dubna, July 1997; Phys. At. Nucl. V61 (1998) 938.
7. ALEPH Collaboration, *Higgs limits in the MSSM: beyond the benchmark*, ALEPH 98-039, CONF 98-018, May 29, 1998.
8. DELPHI Collaboration, *MSSM Parameter Scan at $\sqrt{s} = 183$ GeV*, DELPHI 98-124, CONF 98-185, June 22, 1998.
9. OPAL Collaboration, *A Search for Neutral Higgs Bosons in the MSSM and Models with Two Scalar Field Doublets*, CERN-EP/98-029, Feb. 20, 1998, accepted by Euro. Phys. Journal C.
10. DELPHI Collaboration, *Search for charginos, neutralinos and gravitinos at $\sqrt{s} = 183$ GeV*, DELPHI 98-66 CONF 134, June 22, 1998.
11. P. Chankowski and St. Pokorski, *Supersymmetric loop effects*, in Perspectives on Supersymmetry, Ed. G.L. Kane, World Scientific, Singapore (1997) p. 27.
12. DELPHI Collaboration, *Search for neutral Higgs bosons in e^+e^- collisions at $\sqrt{s} = 183$ GeV*, DELPHI 98-95 CONF 163, June 22, 1998.